

This report is preliminary and has not been reviewed for conformity with U.S. Geological editorial standards or with the North American Stratigraphic Code. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

OPHIOLITIC COMPLEXES AND ASSOCIATED ROCKS IN UST – BELAYA MOUNTAINS AND ALGAN RIDGE, RUSSIAN FAR EAST

compiled by
S.A. Palandzhyan and G.G. Dmitrenko
1996




Fig. 1. TiO_2 versus FeO/MgO variation diagram for mafic volcanics and dikes. OIB – ocean island basalts; MORB – mid-ocean ridge basalts; IAT – island arc tholeites. Fields of different-type basalts from Glassley, 1974.




Fig. 2. K_2O versus SiO_2 diagram for granitoids, after Coleman, 1977. 1 – oceanic plagiogranite; 2 – continental trondjemite.




Fig. 3. Normative plagioclase versus RO/SIO variation diagram for tectonized peridotite, after Palandzhyan, 1992. Fields of average data for diverse types of mantle peridotite: 1 – least depleted sub-continental; 2 – oceanic; 3 – depleted sub-continental; 4 – depleted oceanic; 5 – depleted sub-continental; 6 – oceanic; 7 – depleted oceanic. Groups of mantle peridotites according to tectonic setting of their formation: 1 – sub-continental, 2 – abyssal near passive margins, 3 – very low spreading ridges, 4 – low and fast spreading ridges, 5 – regions of upper mantle with superimposed hydrous partial melting (supra-subduction zone and/or intraplate hot-spots area).




Fig. 4. Na versus Cr in clinopyroxene variation diagram for tectonized peridotite (in oxygen fugacity for 6 oxygen after Komarobat and others, 1981). Fields of average data for diverse types of mantle peridotites according to tectonic setting of their formation: 1 – sub-continental, 2 – abyssal near passive margins, 3 – very low spreading ridges, 4 – low and fast spreading ridges, 5 – regions of upper mantle with superimposed hydrous partial melting (supra-subduction zone and/or intraplate hot-spots area).




Fig. 5. Al_2O_3 in orthopyroxene (opx) versus $Cr/(Cr+Al)$ spinel (sp) variation diagram, after Bonatti and Michael, 1989, revised by Palandzhyan, 1992. Groups of tectonized peridotite according to tectonic setting of their location: 1 – sub-continental, 2 – late Cretaceous and Paleogene; 3 – mid-oceanic ridges, 4 – west-Pacific island arcs inner slope of trench, 5 – very depleted harzburgite of the west Pacific Mesozoic arcs and fragments in Mainits Zone of Koryak Highland, and in Papua, New Guinea.

Fig. 6 Location of the Ust-Belaya subterrane

Fig. 6 Location of the Ust-Belaya subterrane